


INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/68, A61K 39/165	A1	(11) International Publication Number: WO 96/30544 (43) International Publication Date: 3 October 1996 (03.10.96)
(21) International Application Number: PCT/GB96/00706 (22) International Filing Date: 22 March 1996 (22.03.96) (30) Priority Data: 9506298.0 28 March 1995 (28.03.95) GB (71)(72) Applicant and Inventor: WAKEFIELD, Andrew, Jeremy [GB/GB]; 43 Taylor Avenue, Kew Gardens, Surrey TW9 4EB (GB). (74) Agents: STEBBING, Peter, John, Hunter et al.; Ablett & Stebbing, 45 Lancaster Mews, Lancaster Gate, London W2 3QQ (GB).	(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	

(54) Title: DIAGNOSTIC AND THERAPEUTIC SYSTEM FOR CROHN'S DISEASE AND COLITIA ULCEROSA



(57) Abstract

The present invention relates to a diagnostic system for the detection of Crohn's disease and ulcerative colitis by means of nucleotide amplification reactions, *in situ* or *in vitro* which detection means are specific for the measles virus and are capable of distinguishing between the "wild" type and the attenuated measles virus strains. The invention further provides a use in the manufacture of a medicament for the treatment of Crohn's disease and/or ulcerative colitis which medicament comprises a suitable vector carrying antisense RNA to specific viral proteins which are consequently inhibited from being expressed in the host.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

- 1 -

Diagnostic and therapeutic system for Crohn's disease and colitis ulcerosa

The present invention relates to a diagnostic system for the detection of Crohn's Disease and ulcerative colitis and to a
5 therapeutic system derived therefrom. No such diagnostic system is currently available.

Until recently there has been no definite cause attributable to the onset of Crohn's Disease or ulcerative colitis which
10 have remained substantially incurable. The best that medicine can provide is some alleviation of the symptoms.

Crohn's Disease afflicts approximately 4 patients per 100,000 of the population, or in the United Kingdom 2,200 new cases
15 per year. Because in its earlier stages Crohn's Disease tends to present as bowel irritation which is much more common, a reliable diagnostic system is required.

Similarly the incidence of ulcerative colitis is sufficient
20 to warrant early diagnosis to enable early treatment.

The applicant has now shown that the causative agent for Crohn's Disease is the measles virus and accordingly the present invention utilises this finding to provide a
25 diagnostic system for the location of a measles virus in bowel tissue, bowel products or in suitable body fluids such as blood or lymph. Continuing studies also increasingly support the finding that measles virus is the causative agent for ulcerative colitis. The invention also provides the basis for
30 an anti-viral therapeutic system for the measles virus.

According therefore to a first aspect of the present invention, there is provided a diagnostic system for the detection of Crohn's Disease and ulcerative colitis which
35 system comprises means for detecting measles virus RNA or a distinctive metabolic product thereof. Such means may be an antigenic system, or a system utilizing a nucleic acid

- 2 -

amplification or hybridization reaction. In the latter case the invention specifically may provide means for performing a reverse transcription polymerase chain reaction or a nucleic acid sequence based amplification reaction or a ligase chain
5 reaction. The systems may including a buffered primer specific for the reverse transcribed DNA from a RNA measles virus and/or an RNA template. The primer may comprise a 5' modified oligonucleotide sequence specific for any measles virus genomic or antigenomic or messenger RNA. The primer may
10 be attached to a labelling or coagulating moiety such as a fluorochrome for ease of analysis.

The distinctive metabolic product may be selected from a gene sequence or metabolic product thereof to form a nucleiocapsid
15 protein, a phosphoprotein, a large protein, an RNA polymerase complex, a matrix protein, a fusion protein or a haemagglutinin protein, specific for the measles virus or a related paramyxovirus.

20 The nucleotide amplification may be a reverse transcription-polymerase chain reaction (RT-PCR) or nucleic acid sequence based amplification NASBA (or 3SR). In the former case a kit for performing the diagnostic tests as hereinbefore set forth may comprise:-

- 25 (1) M-MLV reverse transcriptase;
(2) random hexamers and/or oligo (Dt)₁₂₋₁₈;
(3) a reaction buffer for (1) above;
(4) a PCR reaction buffer including Taq DNA polymerase
(5) a 5' modified PCR primer with reporter molecules.

30

Such a kit may further comprise a positive control which comprises measles virus RNA in solution at a known concentration and preferably a negative control or means for its provision.

35

- 3 -

We have however now found that NASBA is the preferred diagnostic method because it is significantly more sensitive for the detection of the measles virus.

- 5 According to a further aspect of the present invention there is provided a method for the *in situ* analysis of a tissue for Crohn's Disease or ulcerative colitis which method comprises the steps of:-
- a) obtaining a tissue sample and securing the same in an
10 enclosed reaction container;
 - b) adding a reagent (1) comprising a PCR buffer; $MgCl_2$, DNTP'S, random hexamers and diluting to a desired dilution factor with an aqueous solution;
 - c) subsequently closing the enclosed reaction container and
15 adding M-MLV-RT and subjecting to heat cycling for at least one cycle;
 - d) treating with a washing buffer;
 - e) adding a buffered Taq DNA polymerase, $MgCl_2$, DNTP and a primer at a predetermined dilution in sterile distilled water
20 and then heat cycling for at least 15 cycles by single or nested PCR reactions; and
 - f) subsequently repeating the steps and viewing for labelled product to indicate the presence of a measles virus RNA.
- 25 In a further aspect of the present invention, there is provided a method for the *in vitro* analysis of a sample for Crohn's Disease or ulcerative colitis which method comprises the steps of extracting measles virus RNA from a tissue sample and
- a) adding thereto a buffer further comprising DTT, DNTP'S
30 and R Nase inhibitor and oligo (dT)₁₂₋₁₈ and letting down to a desired aqueous concentration in an enclosed reaction container;
 - b) adding thereto M-MLV-RT and subjecting to incubation to
35 provide a cDNA product;
 - c) purifying the product and adding thereto a PCR buffer along with $MgCl_2$, dNTP's and an outer primer;

- 4 -

- d) subsequently adding Taq DNA polymerase and heating for at least 15 cycles by single or nested PCR reactions;
- e) recovering an aliquot of a so-formed reaction production and adding buffered PCR reaction mixture with an inner pair of primers and further Taq DNA polymerase and heat cycling for at least 20 cycles; and
- f) moving the so-formed product and adding a loading dye and subjecting the resultant product to an electrophoresis to identify a resultant product band.
- g) the sequencing of the amplified products, or hybridising the amplified products with a homologous or heterologous specific probe in order for example to distinguish vaccine strain measles virus from "wild" type.
- According to a further aspect of the present invention there is provided a medicament for the treatment of Crohn's disease and/or ulcerative colitis which method prevents expression, replication, transcription, RNA processing and/or mRNA transport of the measles virus in the host. The medicament may include a measles virus antisense RNA for either genomic or antigenomic RNA and a vector.

The medicament may comprise a measles virus peptide or carbohydrate antigen or a monoclonal or polyclonal analogue thereof. The invention also comprises a measles virus-specific nucleic acid based vaccine genome encoded into an expression system adapted to induce a specific immune response to the measles virus and/or its component antigens.

Using antisense RNA to the viral RNA including messenger RNA for the treatment of measles associated diseases alleviates the problem of side effects on host cellular functions. In the past such treatments have proved elusive partially due to the fact that it is extremely difficult to design inhibitors specific for the essential proteins of a virus. However, now in the prior art, the antisense RNA approach has been successfully used to inhibit viral genes of avian retrovirus,

- 5 -

Rous sarcoma virus, human immunodeficiency virus and simian immunodeficiency virus in cultured cells.

The invention provides a medication which comprises a suitable
5 vector for uptake by the infected cells engineered to carry
the appropriate measles virus antisense RNA designed to be
targeted against the appropriate gene (e.g Nucleoprotein gene
and/or Haemagglutinin gene) followed by a drug resistance gene
(e.g Hygromycin B). The method of preparation of the
10 medication is a modification of the methods in Koschel K. et
al (1995), Virology, 207, 168-178.

The invention will now be described by way of illustration
only in the following examples and with reference to the
15 attached Figures wherein:-

Figure 1a shows

Measles infected Vero cells immunostained with measles primary
polyclonal antibody: areas of syncytial cytopathic effect
20 exhibit intense cytoplasmic staining. Counterstained with
Mayer's haematoxylin (original magnification x 350);

Figure 1b shows

Immunogold labelled measles infected Vero cells. Frequent
25 foci of double and often larger clusters of gold particles are
seen, associated with typical fibrillar arrays of
nucleocapsids, 15-20 nm in diameter. Counterstained with
uranyl acetate and Reynold's lead citrate (original
magnification x 91,000);

30

Figure 1c shows

Measles infected Vero cells treated identically to 1a, but
omitting the measles antibody primary (original magnification
x 375);

35

- 6 -

Figure 1d shows

Measles infected Vero cells developed without measles primary antibody, but exposed to gold-conjugated secondary antibody; no labelling is seen (original magnification x 91,000);

5

Figure 1e shows

Mumps primary antibody on measles infected Vero cells, showing no signal (immunoperoxidase; original magnification x 400).

10

Figure 2a shows

Mumps infected Vero cells immunostained with mumps primary antibody showing cytoplasmic staining in infected cells (original magnification x 400);

15

Figure 2b shows

Mumps infected Vero cells treated identically to 2a, but omitting the mumps primary antibody (immunoperoxidase; original magnification x 400);

20

Figures 2c and 2d show

Measles primary antibody on mumps infected Vero cells examined by immunoperoxidase (2c) (original magnification x 400) and immunogold (2d) (original magnification x 82,400). No antibody cross reactivity is seen. In 2d, the fine fibrillary
25 arrays of mumps nucleocapsids are seen clearly;

Figure 3 shows

SSPE: immunostaining for measles virus nucleocapsid protein in SSPE brain examined by light microscopy (Figure 3a)

30 original magnification x 400) and electron microscopy (Figures 3b and 3c). Intranuclear nucleocapsids are seen in Figure 3b, frequently showing paired gold particles spanning individual nucleocapsids, demonstrated clearly in Figure 3c, showing cytoplasmic signal (original magnifications; 3b x 62,500, 3c
35 x 95,000). Figure 3c, showing cytoplasmic signal (original magnifications; 3b x 62,500, 3c x 95,000). No signal is seen in identically processed tissue sections, in which the primary

- 7 -

measles antibody was omitted (Figure 3d; immunoperoxidase original magnification x 400);

Figure 4 shows

5 Measles appendicitis: immunoperoxidase treated sections stained for measles virus N protein. Positive signal is seen in an endothelial cell syncytium (Figure 4a) (erythrocytes arrowed) (original magnification x 100) and a Warthin Finkeldey giant cell (Figure 4b) (original magnification x
10 800);

Figure 5 shows

Crohn's Disease: positive nuclear staining in macrophage-like cells in a focus of granulomatous inflammation (Figure 5a; immunoperoxidase; original magnification x 600). Signal is
15 not seen in an identically processed serial section, omitting the primary antibody (Figure 5b, original magnification x 600). Positive immunogold labelling is seen in an identical cellular location and shows the characteristic paired signal
20 (arrowed) Figure 5c, original magnification x 71,000);

Figure 5d (inset) shows a high power view of gold label distributed along the length of what appears to be a single nucleocapsid (original magnification x 139,000);

25

Figure 5e shows nuclear staining in a macrophage in a focus of granulomatous inflammation from another case of Crohn's Disease. The paired signal is observed, particularly in the central cluster, shown magnified in 5f; this demonstrates
30 clearly, the parallel arrays (arrowed) of a 17 nm diameter nucleocapsid spanned by the gold label (original magnifications; Figure 53 x 91,000, Figure 5g x 95,000). Staining was not observed in otherwise identically processed sections of Crohn's Disease when the measles primary antibody
35 was omitted (Figure 5g, original magnification x 54,000).

EXAMPLE 1

A study was conducted to confirm the viral origins of the nucleocapsid-like particles observed in bioassays of bowel
5 tissue in Crohn's Disease or probably ulcerative colitis by co-localising these particles with a gold-labelled antibody specific for measles virus nucleocapsid protein. These particles have not previously been detected ultrastructurally in normal intestine.

Cell Culture

Vero (green monkey kidney) cells were cultured in 75 mm flasks using standard techniques. Confluent monolayers were infected with either Edmonston strain measles virus at a titre of 10^7
15 plaque forming units per ml, or mumps virus (Taylor strain: PHLS Colindale, UK) at a TCID₅₀ (Tissue Culture Infectious Dose) of 10^4 ; uninfected cells were used as a negative control. When > 80% of cells showed cytopathic effect, monolayers were fixed using 1% glutaraldehyde/1%
20 paraformaldehyde in phosphate buffered saline (PBS) for 30 minutes. The cells were scraped from the flask using platinum wire loop and the suspended cells were pelleted in Eppendorf tubes using a Beckman microfuge (Palo Alto, California, USA). Cell pellets were washed after fixation with PBS and
25 dehydrated using dimethylformamide, 50, 70 and 90% x 2 changes each of 10-20 minutes duration depending on the size of the block. The blocks were then infiltrated with a 50:50 mixture of dimethylformamide and LR White resin with 0.5% benzoin photoinitiator (TAAB Laboratories Equipment Ltd, Reading, UK)
30 added, for 30-60 minutes followed by 100% resin for 1-2 hours. The blocks were embedded in closable embedding capsules using fresh LR White with photoinitiator. Polymerisation was carried out at 4° using a UV light source at 10cm distance. The resin polymerised in 1-2 hours and the small amount of
35 resin remaining unpolymerised under the closure was removed using a cotton swab.

- 9 -

Tissue Reprocessed from Paraffin Blocks

Initially, archival formalin-fixed paraffin processed tissues with established measles infection were selected; these included cerebral tissue from a case of subacute sclerosing pan-encephalitis (SSPE), and appendix from a case of acute measles virus appendicitis (Gift of Dr H Reid, Chase Farm Hospital, Enfield). Six cases of granulomatous Crohn's Disease were selected that included 4 rectal biopsies and two appendices taken at the time of initial presentation (that is, before specific or immunosuppressive therapy was instituted). In both the positive controls and Crohn's Disease tissues, the diagnoses were established by standard clinical and histopathological criteria. Two cases of ileocecal tuberculosis (TB) were studied as granulomatous controls. Sections were cut from tissue blocks and immunostained with a polyclonal antibody specific for the measles virus nucleocapsid (N) protein (gift of J Stephenson and T Fookes, CAMR, Porton Down, UK) using an immunoperoxidase technique, as described previously (Wakefield AJ. J Med Virol 1993; 39:345-353). Serial tissue sections were processed with either omission of the primary antibody or exposure to mumps (a related paramyxovirus) specific monoclonal antibody (Seralab, Crawley, Sussex, UK) as negative controls. Cross-reactivity of the measles and mumps specific antibodies was examined in Vero cells infected with these viruses, prior to immunogold studies.

Processing for Electron Microscopy

Suitable areas of tissue were selected for immunogold analysis; these included areas that gave positive measles staining by immunoperoxidase, and foci of granulomatous inflammation in the cases of Crohn's Disease and ileocecal TB. Wedges of tissue were removed from the paraffin block using a single edged razor blade. The pieces were dewaxed with chloroform followed by rinsing in absolute alcohol. The tissue was then infiltrated with resin and embedded as described above.

- 10 -

Paraffin sections only were available on the acute measles
appendicitis. A novel method was used to lift the tissue from
the slide; the section was dewaxed and taken to absolute
alcohol, it was then flooded with 50:50 mixture of alcohol and
5 LR White resin with photoinitiator added for a further 15
minutes; it was then drained and the slide blotted to remove
surplus resin. Drops of fresh resin were placed over the
tissue and covered with plastic coverslips (Agar Scientific,
Stansted, UK). The resin was then polymerised with UV at 4°C
10 for one hour. The coverslip was peeled off and unpolymerised
resin removed. The resin containing the tissue was lifted by
immersing the slide in liquid nitrogen for a few seconds. The
pieces were then trimmed and mounted onto blank resin blocks
using Permabond quick setting epoxy resin (Permabond Adhesives
15 Ltd, Eastleigh, UK) Ultrathin sections were cut at 50-80 nm
and immunostained as above.

Immunolabelling

Ultrathin section were cut at 70-80 nm and picked up onto
20 PiliForm (Agar Scientific, Stansted, UK) coated 300 mesh
nickel grids (Gilder, Grantham, UK). The grids were incubated
on drops of 5% normal goat serum in 0.1% BSA/PBS buffer for
30 minutes, they were then transferred to drops of primary -
measles polyclonal antibody 1/100 in 0.1% BSA/PBS for 1 hour.
25 Primary antibody was removed by washing on drops of PBS 5 x
5 minutes, the first transfer being made without removing any
surplus liquid from the grid. Surplus buffer was removed on
the remaining four changes by touching the edge on a piece of
blotting paper, taking care not to allow the grid to dry out.
30 The grids were then transferred to drops of gold conjugate
(Biocell, Cardiff, UK) diluted 1/100 with PBS for 1 hour
followed by washing with drops of PBS 2 x 5 minutes to remove
unbound gold conjugate and a final rinse in distilled water.
Sections were stained lightly with uranyl acetate and
35 Reynold's lead citrate and viewed using a Philips 201
transmission electron microscope. Sections processed omitting

- 11 -

the primary antibody, were included with all tissues as negative controls.

Measles-infected Vero Cells

- 5 Vero cells that had been experimentally infected with measles virus exhibited positive immunoperoxidase and immunogold labelling (Figures 1a and 1b) that was not seen either in identically processed measles infected cells not exposed to the primary measles antibody (Figure 1c and 1d), or in measles
10 infected cells exposed to the mumps primary antibody (Figure 1e). Ultrastructurally, viral nucleocapsids consisted of characteristic parallel fibrillar structures of between 15 and 20 nm diameter.
- 15 A characteristic pattern of labelling was observed in both nucleus and cytoplasm of infected cells that consisted of gold particles - grouped, often in pairs, and bound at discrete points along the viral nucleocapsids (Figure 1b); this pattern was seen consistently in both infected cells and infected
20 tissues, and was not seen in the very occasional background signal observed in control sections. Mumps infected Vero cells stained strongly by immunoperoxidase using the primary mumps antibody (Figure 2a), but did not stain when the mumps primary antibody was omitted in otherwise identically
25 processed cells. No signal was observed when the measles antibody was put onto mumps infected cells at either the light microscopic or ultrastructural levels.

Subacute Sclerosing Panencephalitis

- 30 Strong signal for measles virus as observed predominantly in the nuclei of infected cells, using both the immunoperoxidase technique on histological sections, and immunogold labelling of ultrathin sections of infected brain. Ultrastructurally, viral nucleocapsids were well preserved despite suboptimal
35 fixation and paraffin processing. Again, the double immunogold signal on nucleocapsids was a consistent feature. No signal was observed on sections in which the primary

- 12 -

measles antibody had been omitted, either at the light microscopic (Figure 3d) or ultrastructural levels (data not shown).

5 Intestinal Tissues

The techniques were applied subsequently to measles-infected intestinal tissue - acute infected measles appendicitis - with similar results. Measles virus antigen was identified by the immunoperoxidase technique in endothelial cells, and
10 macrophage-like cells including Warthin-Finkeldy giant cells, a classical feature of acute measles virus infection, and occasional lymphocytes within lymphoid follicles. Staining was not seen in negative control sections. Immungold labelling confirmed the presence of the virus in the same
15 cellular foci despite suboptimal preservation. No signal was observed in the absence of the primary antibody.

Of the six cases of Crohn's Disease examined, all were positive for measles by immunoperoxidase within cells
20 consistent in morphology and location with histiocytic macrophages, endothelial cells and occasional lymphocytes; signal was not seen in sections that were either not exposed to the measles primary antibody, or exposed to mumps primary antibody. Five cases were positive by immunogold in the same
25 cellular location, and viral particles exhibited the characteristic features of size, shape and immunogold labelling described above; in one case, signal was detected additionally, in a single crypt epithelial cell. In the fifth case, that was negative by immunogold, the focus of
30 granulomatous inflammation present in the histological sections had "cut out" and could not be identified in the ultrathin section. Crohn's sections, processed in an identical manner, but excluding the measles primary antibody, were negative. Of the two cases of ileocecal TB that were
35 examined, neither was positive by immunoperoxidase, although one case exhibited a low level of nuclear signal in a macrophage-like cell, by immunogold labelling.

- 13 -

This presents the first direct evidence of measles virus persistence in the intestine; that is, the co-localisation of a specific antibody with viral nucleocapsids at the ultrastructural level. The cellular localisation of measles virus in Crohn's tissues is consistent with that observed previously using different techniques (Wakefield AJ J Med Virol 1993; 39; 345-353; Knibbs DR, Gastroenterology, 1993; 104; A726 (Abstract)), and in different laboratories (Knibbs DR. Gastroenterology, 1993;104;A726 (Abstract)). The co-localisation of a specific antibody with particles that are, in terms of both size and morphology, identical to the target virus, is strong evidence for the presence of that virus; the characteristic pattern of labelling of nucleocapsids with gold particles provided a further, if unexpected degree of specificity. The detection of measles in one of two cases of ileocecal TB raises the possibility that persistently infected immune cells aggregate in foci of inflammation, and are unrelated to the primary cause of the granuloma. This notwithstanding, in Crohn's Disease tissues, the detection of measles virus within resident intestinal cells, including endothelium and epithelium, does suggest persistent intestinal infection. It is possible however that circulating immune cells within the same host may also be persistently infected.

Corticosteroids, a mainstay in the treatment of patients with Crohn's Disease, may predispose to permissive viral replication, and confound the question of persistence within the inflamed intestine; none of the patients whose tissues were examined in this study had received either corticosteroids or other immunosuppressive therapy.

EXAMPLE 2

The implied aetiological association of measles virus with Crohn's Disease is supported by detection of an immune response to infected cells in affected tissues. This example sought to detect and characterise *in situ* immune responses to

- 14 -

measles virus in both acutely and persistently infected tissues, and in particular, Crohn's granulomata. Serial tissue sections from cases of Crohn's Disease (n = 17), tuberculosis (n = 9), acute intestinal ischaemia (n = 5), acute measles pneumonitis (n = 2), acute measles appendicitis (n = 1), subacute sclerosing panencephalitis (SSPE; n = 1), and measles inclusion body encephalitis (MIBE; n = 1), were examined. Single and double immunohistochemical labelling was performed to identify both cytotoxic lymphocytes (CD8, TIA, perforin, Leu 7, CD45RO, CD45RA) and macrophages (KP1). The relationship of these cells to measles infected cells was identified by double immunolabelling with anti-measles virus nucleoprotein antibody. In both acute measles appendicitis and SSPE, CD8⁺/TIA⁺ cytotoxic lymphocytes (CTL) targeted infected cells. In the other tissues that were positive for measles virus including Crohn's Disease (13/17) - where staining was largely confined to granulomata, MIBE, fatal pneumonitis, and 1 tuberculous granuloma, infected cells appeared to be targeted by macrophages rather CTL. The CTL in Crohn's granulomata were Leu 7⁻ and perforin⁻/CD45RO⁻ (naive). CTL in both tuberculous and Crohn's granulomata were similar in their peripheral distribution, number and phenotype. The data suggest that measles-specific CTL responses may be attenuated in Crohn's Disease compared with acute measles appendicitis and SSPE, and secondly, that an abnormal macrophage response to persistent measles virus infection of the intestine could result in granulomatous inflammation.

30 EXAMPLE 3

In an international collaborative study, serum measles IgM immunoreactivity was assayed by ELISA in consecutive outpatients with Crohn's Disease (n = 95), ulcerative colitis (n = 79), viral hepatitis (n = 63) and blood donors (n = 30). Results were compared with those obtained from a different commercial measles IgM assay, serum rubella and Epstein-Barr

- 15 -

virus-specific IgM immunoreactivity, total serum IgM, Rheumatoid Factor and measles-specific IgG. Twenty patients with inflammatory bowel disease were studied serially over a four month period. At the ELISA cut-off point for
5 confirmation of recent acute measles virus infection, there were no significant differences between groups. However, increased serum measles IgM immunoreactivity (\geq mean \pm 2SD of blood donors) was significantly greater in patients with Crohn's Disease 39/95 (41%) and ulcerative colitis 33/79 (42%)
10 compared with hepatitis patients 5/64 (8%) and normal controls 0/30 (0%) ($p < 0.001$). Those positive by ELISA were also positive by indirect immunofluorescence on the same serum sample. Serum measles IgM immunoreactivity did not correlate with either total IgM, rubella or Epstein-Barr virus IgM (not
15 raised), measles IgG, or disease activity. Patients not receiving steroids were more likely to have raised measles IgM immunoreactivity ($p < 0.05$). All sera examined for Rheumatoid Factor were negative. Of twenty patients with inflammatory bowel disease studied over four months, 55%
20 showed raised measles IgM immunoreactivity at some stage during the follow-up. The data suggested an immunological response to measles virus in patients with Crohn's Disease and ulcerative colitis, supporting a potential aetiological role for this agent in inflammatory bowel disease.

25

EXAMPLE 4

The epidemiological association between Crohn's Disease and early measles virus exposure has been indicated either
30 indirectly, on in studies of case-control design. In order to determine absolute risk estimates for in utero measles exposure and Crohn's Disease, maternity charts for all 25,000 deliveries at University Hospital, Uppsala, between 1940-1949 were reviewed; four cases of overt measles infection in the
35 mother during pregnancy were identified. Offspring, and in two cases their mothers also, were interviewed, and case records reviewed. Cases, 1, 2 and 3 had undergone multiple

- 16 -

intestinal resections; tissues, available from two cases, were examined by both routine histology, and immunohistochemistry and immunogold electronmicroscopy for measles virus, using appropriate positive (brain; subacute sclerosing panencephalitis) and negative (no primary antibody, mumps infected cells) controls (Gut. 1995; 36: 564-9). Three cases of Crohn's Disease occurred in the four offspring; in each case the disease was preceded by recurrent, antibiotic resistant pneumonia. All had extensive ileal and colonic disease, two patients requiring intravenous feeding. Of the four cases, the only one to have suffered clinical acute measles disease, did not develop Crohn's Disease. Tissues from two cases of Crohn's Disease that were examined for measles virus antigen, were positive in foci of granulomatous and lymphocytic inflammation. The data indicate that in utero exposure to measles virus is a major risk factor for severe, extensive Crohn's Disease. Exposure at this time may lead to persistent infection, or alternatively, modify the response to the infection in later life, leading to virus persistence.

EXAMPLE 5

Immunogold electron microscopy was used to test for the presence of measles virus in intestinal tissue from patients with Crohn's Disease, and inflammatory and non-inflammatory disease controls. Formalin-fixed, paraffin-embedded tissue was reprocessed and stained with anti-measles nucleoprotein primary antibody followed by 10 nm gold-conjugated secondary antibody. Similarly processed brain tissue, taken from a patient with subacute sclerosing panencephalitis (SSPE), was used as the positive control. Duplicate sections of all tissues were processed without the primary antibody. In Crohn's Disease 8/9 foci of granulomatous inflammation and 0/4 foci of non-specific inflammation were positive for measles virus. Of control,s 0/5 ulcerative colitis tissues and 1/1 SSPE tissues were positive. Gold grain counts per nuclear field in both Crohn's granulomata and SSPE were highly

- 17 -

significantly greater than controls, including non-granulomatous areas of Crohn's Disease ($p < 0.0006$); in both diseases staining was confined to a small population of cells exhibiting characteristic cytopathology. These data support
5 a role for measles virus in the aetiology of Crohn's Disease.

EXAMPLE 6

Kits for Measles Virus Detection

10 Measles is an RNA virus. The nucleic acid amplification mediated detection whether *in situ* or *in vitro* of this example is based on either reverse transcription - polymerase chain reaction (RT-PCR) or nucleic acid sequence based amplification NASBA (3SR).

15

The main components of the RT-PCR kit for measles detection *in vitro* or *in situ* possess

- (i) Reverse transcriptase eg M-MLV reverse transcriptase
- 20 (ii) Primers for first strand cDNA synthesis
- iii) reaction buffer for RT,
- (iv) PCR reaction buffer,
- (v) PCR primers which may be 5' modified with reporter molecules, eg fluorochromes.

25

Additionally an Amicon Microcon 30 size exclusion cartridge is optionally required for purification of the product from first strand DNA synthesis for *in vitro* amplifications.

30 Both kits preferably include a positive control, for example measles virus RNA, supplied as a solution at known concentration with which individuals using the kit can calibrate their results.

35 Additionally a negative control is required. This is preferably achieved by treatment of one portion of the sample to be tested with RNase A for 10 minutes at 37°C prior to

- 18 -

analysis. This is effective for both *in situ* and *in vitro* applications and additionally indicates any non-specific product formation from any DNA that may be present as a contaminant in the samples. It is therefore desirable to
5 include Rnase A as an additional component in the kit.

Two basic protocols follow for a) *in situ*; and b) *in vitro* amplification.

10 EXAMPLE 7

a) Measles virus detection *in situ* - Kit 1

1) To the tissue section as prepared in Example 6 is added
15 75 μ l of reagent (1). The slide is sealed in an enclosed reaction chamber such as that shown in PCT/GB95/00215 and placed on a flat block of a pHC3 Techne thermal cycler. With the enclosed reaction container so positioned 200 units of M-MLV RT are added and heated for 2 hours at 42°C, followed by
20 five minutes at 95°C, followed by five minutes at 15°C to complete a first heating cycle.

2) The slide is then thoroughly rinsed in the reagent (2),
and

25

3) Subsequently the tissue section is covered with 430 μ l of the reagent (3). To this is added 1.25 units of Taq DNA polymerase and the whole is heated for five minutes at 95°C followed by thirty heat cycles of two minutes at 58°C, 1.5
30 minutes at 75°C and 1 minute at 95°C. The action is finished by heating for ten minutes at 72°C.

4) The slide is then rinsed thoroughly in reagent (2) and viewed under an epifluoresence microscope.

35

- 19 -

Reagents for Kit 1 in situ measles detectionReagent 1 (RT buffer - inc. primers)

- q
5 10 x PCR buffer
MgCL₂
dNTPs
RNA Guard
Random Hexamers
10 DEPC treated water

At required final
working concentration

- 15
Reagent 2 (Washing buffer)
phosphate buffered saline Ph 7.5

Reagent 3 (PCR buffer - inc. primers)

- 20 10 x PCR buffer
MGCL₂
DNTPS
Primer 1) either 5' modified
Primer 2) or +Dig-UTP
25 Sterile double distilled water

At required final
working concentration

30

Enzymes

- M-MLV RT - (Eg Gibco BRL)
35 Taq DNA polymerase - (eg Gibco BRL)

EXAMPLE 8

40

- b) Measles nucleic acid detection by amplification in vitro
Kit 2

It is assumed that total RNA will have been prepared from
45 tissues of interest for example by the method of Chirgwin et
al (1979) Biochemistry, 18, 5294-5299, the diagnostic test may
then be effected by the following steps:

- 20 -

- 1) To 0.4 μ g of RNA add sufficient 5 x reagent (4) 1 and DEPC treated sterile double distilled water to make up volume to 20 μ l.
- 5 2) Remove an aliquot (containing 100 ng) of total RNA for an internal control reverse transcriptive PCR using the low copy number cellular UIA RNA as a target for amplification;
- 10 3) Perform hybrid capture on UIA positive RNA samples using magnetic beads with amino-linked oligonucleotide specific for both positive and negative measles RNA strands;
- 4) Separate the magnetic beads and elute measles RNA with 50 μ l of elution buffer;
- 15 5) Use 5 μ l of eluent for measles reverse transcription PCR together with relevant positive and negative controls.
- 20 6) Add 200 units of M-MLV RT and incubate at 42°C for 2 hours for 30 minutes
- 7) Take the resultant cDNA product and purify using an Amicon Microcon 30 cartridge according to the manufacturers instructions.
- 25 8) Take 10 μ l of purified cDNA product from step 3 and make up to 39 μ l with a total of 10 x reagent (5) and sterile double distilled water.
- 30 9) Add 1 unit of Taq DNA polymerase and heat for 10 minutes at 95°C followed by 35 heat cycles of 58°C for one minute, 72°C for 1 minute and 95°C for 1 minute. Finished with one heat cycle of 72°C for 5 minutes
- 35 10) Take 1 μ l of the reaction product from step 5 and make up to 23 μ l with 10 x reagent (6) and sterile double distilled water.

- 21 -

11) Add 1 unit of Taq DNA polymerase and heat cycle as for step 5 but for 25 rather than 35 cycles.

12) Take 18 μ l of product from step 7 add 2 μ l of loading dye and electrophorese on a 1% agarose gel to identify product of a desired kbp.

Reagents for Kit 2 in vitro measles detection

10	<u>Reagents 4 (RT buffer)</u>	<u>Reagent 5</u>	(PCR buffer 1) - inc primers	<u>Reagent 6</u>	(PCR buffer 2) - inc primers
	RT buffer	PCR buffer		PCR buffer	
	DTT	MgCL ₂		MgCL ₂	
	dNTPs	W - 1		W - 1	
	RNase inhibitor	dNTPs		dNTPs	
15	Oligo (dT) ₁₂₋₁₈	MV1)			
			outer primers		inner primers
		MV2)		MV4)	
20	At 5 x working concentration	At 10 x working concentration		At 10 x working concentration	

Enzymes

M-MLV RT - (eg Gibco BRL)
25 Taq DNA polymerase - (eg Gibco BRL)

Additionally

DEPC treated sterile double distilled water
30 Amicon Microcon 30 size exclusion cartridge.

EXAMPLE 9

NASBA amplification of measles virus N gene sequence

1. Warm NASBA buffer and primer, mix at room temperature and make up master mix for 20 μ l reactions in clean room.
2. Aliquot master mix into Eppendorf tubes and add 100 ng of template RNA per reaction prepared by the method of Cosby et al 1989.

- 22 -

3. Maintain tubes at 65°C for 5 minutes to denature RNA then hold at 41°C for 5 minutes.
4. Warm enzyme mix to room temperature then add to reaction mix,
- 5 5. Incubate at 41°C for 90 minutes in a water bath.
6. Detect reaction products via Northern blotting or use an enzyme-linked gel assay (e.g ELGA, Organon or Technika).

NASBA BUFFER

10	REAGENT	VOLUME (μ l)	FINAL CONCENTRATION
	1M Tris pH8.5	200	40 mM
	1M MgCl ₂	60	12 mM
	4M KCl	87.5	70 mM
15	1M DTT	25	5 mM
	20 mM each dNTP	50 each	1 mM each
	20 mM ATP, UTP, CTP	100 each	2 mM each
	20 mM GTP	75	1.5 mM
	100 mM ITP	25	0.5 mM
20	H ₂ O	27.5	
		1000 μ l	

Aliquot and stored at -20°C

25

PRIMER MIX

	REAGENT	VOLUME (μ l)	FINAL CONCENTRATION
	100% DMSO	750	75%
	50 μ M Primer 1	20	1 μ M
30	50 μ M Primer 2	20	1 μ M
	H ₂ O	1000 μ l	

Aliquot and store at -20°C

- 23 -

ENZYME MIX

REAGENT	VOLUME (μ l)	FINAL CONCENTRATION
H ₂ O	969	1.5 M
5 4.5 M Sorbitol	668	105 ng/ μ l
20 mg/ml BSA	42	105 ng/ μ l
0.87 U/ μ l Rnase 11	36	0.004 U/ μ l
70 U/ μ l T7 RNA Pol.	184	1.6 U/ μ l
25.3 U/ μ l AMV RT	101	0.32 U/ μ l
10	2000 μ l	

Aliquot and store at -70°C

NB. Do not vortex the enzyme mix.

15 EXAMPLE 10

Measles RNA extraction and detection using NASBA was effected using the following steps:-

- 20 1. Homogenise up to 500 mg of tissue in 3 ml total RNA isolation reagent (Advanced Biotechnologies). Total RNA extraction method is based on the single-step method of RNA isolation by Chomczynski et al. If using cells, resuspend pellet in 1 ml of reagent. If using blood, mix with 5 x vol.
- 25 of reagent.
2. Extract total RNA and resuspend pellet in 100 μ l of DEPC treated dH₂O.
- 30 3. Measure optical density of extracted RNA solution and determine the RNA integrity via gel electrophoresis on a 1.5% agarose gel.

- 24 -

4. Remove an aliquot (containing 100 ng) of total RNA for an internal control NASBA using the low copy number cellular U1A RNA as a target for amplification.
5. Perform hybrid capture on U1A positive RNA samples using magnetic beads coated with amino-linked oligonucleotides specific for both positive and negative measles RNA strands.
6. Separate the magnetic beads and elute measles RNA with 50 μ l of elution buffer.
7. Use 5 μ l of eluent for measles NASBA together with relevant positive and negative controls.
- 15 Detect specific NASBA products using Enzyme Linked Gel Electrophoresis with an internal horseradish peroxidase-labelled oligonucleotide.

Measles NASBA

- 20 • Measles NASBA should be performed using primers specific to the nucleoprotein region of the measles virus genome for example, primers AB20 and AB22.
- Specific NASBA products should be detected using an internal oligo probe for example AB20.

OLIGO	POSITION	SEQUENCE
AB20 (NASBA) (Upstream)	1200 - 1219	AGG GCA AGA GAT GGT AAG GA
AB22 (NASBA) (Downstream)	1358 - 1379	AAT TCT AAT ACG ACT CAC TAT AGG G GA TCA CCG TGT AGA AAT GAC A
AB10 (PROBE)	1288 - 1308	GTT TCA GAG ATT GCA ATG CA

- 25 -

EXAMPLE 11*In situ hybridisation detection of measles virus*

It is possible to detect measles virus RNA *in situ* in tissue
5 sections which have been , for example, paraffin or araldite
embedded or frozen, via *in situ* hybridisation.

Detection of hybrids can be either direct i.e. by fluorescence
or autoradiography or indirect i.e. by subsequent reaction of
10 the hybrid with a reporter molecule to allow its detection,
for example, by chemiluminescence or fluorescence.

The following example method is for the identification of
measles virus RNA in tissue sections employing a biotinylated
15 single stranded RNA probe and the subsequent immuno-detection
of hybrids. This probe is devised from the N gene sequence
(Cosby et al 1989) and is 186 bp in length. It is specific
for all nucleocapsid sequences of measles virus contained in
the GenBank sequence data base but will not react with the
20 closely related morbillivirus - canine distemper virus.

HYBRIDISATION

1. Take either semi-thin or ultra-thin sections and, if
paraffin embedded, dewax and rehydrate (on slide), add 200 μ l
25 of proteinase K (1mg ml⁻¹), and incubate for 15 ml at 37°C.
2. Wash in DEPC treated water and paraformaldehyde fix for
5 min.
3. Wash sections in DEPC treated water.
4. Add hybridisation buffer and incubate at 42°C for 16
30 hours.
5. Wash sections in 0.6% NaCl, 10 mM HCl pH 7.0, Immol EDTA
for 5 minutes at 25°C.
6. Rinse sections in DEPC treated water.
7. Wash sections in 45% v/v formamide in washing buffer for
35 30 minutes at 28°C.
8. Rinse sections in DEPC treated water.
9. Wash sections in 0.1 x SSC for 60 minutes at 40°C.

- 26 -

10. Wash sections in 10mmol phosphate buffered saline twice for 5 minutes each at 25°C.

Solution

5 Hybridisation buffer (Final Concentration)

Formamide (50% v/v)

5 x SSC

5 x Denhardtts solution

0.25 mg ml⁻¹ Salmon Sperm DNA

10 0.5 mg ml⁻¹ yeast tRNA

10% Dextran sulphate

DEPC treated water

Probe RNA \leq 100 nmol

15

DETECTION (adapted from Cosby et al 1989 and McQuaid et al 1990)

1. Monoclonal mouse anti-biotin antibody at 1:40 dilution is applied at 37°C for 30 minutes to the section.

20 2. Wash sections twice in excess phosphate buffered saline.

3. Add biotinylated anti-mouse antibody at a 1:90 dilution and incubate for 30 minutes (ibid) at 37°C.

4. Wash sections twice in excess phosphate buffered saline.

5. Add a 1:500 dilution of peroxidase conjugate and incubate
25 for 30 min at 25°C.

6. Rinse the sections in excess phosphate buffered saline for 8 minutes

7. Add substrate 3-amino-9-diethylcarbazole and leave at 25°C for 10 minutes.

30 8. Wash sections in running water for 10 minutes.

9. Allow to dry and view under microscope.

10. Sections can be counter stained with Mayers haematoxylin if appropriate.

35 A DETECTION KIT using the above technique comprises:-

(a) the riboprobe;

(b) hybridisation buffer;

- 27 -

- (c) proteinase K;
- (d) Rnase A; and
- (e) antibodies and detection reagents.

5 A positive and negative control are included where the negative control may be generated by the pre-treatment of an arbitrary, duplicated sample, with RNase A for 1 hour at 37°C prior to hybridisation and detection. A positive control for the kit may be a sample of plasmid borne transcription
10 template and for section detection the control could be a sample of a known SSPE infected tissue.

The kits in accordance with the present invention locate measles RNA which is indicative of Crohn's Disease from tissue
15 samples from biopsies of the bowel and from body fluids such as blood and lymph and from faecal extracts.

The invention relates therefore to a diagnostic system for the detection of Crohn's Disease and ulcerative colitis to assay
20 kits for utilisation of said system, and to therapeutic systems derived therefrom.

- 28 -

CLAIMS:

1. A diagnostic system for the detection of Crohn's Disease and/or ulcerative colitis which system comprises means for
5 detecting measles virus RNA, or distinctive metabolic product thereof in bowel tissue, bowel products, or a body fluid.
2. A system according to claim 1 wherein said means for detecting virus RNA comprises means for performing a nucleic
10 acid based test either *in situ* or *in vitro*.
3. A system according to claim 2 wherein the test comprises a nucleotide amplification or hybridization reaction selected from a polymerase chain reaction (RT-PCR) or a nucleic acid
15 based amplification reaction (NASBA).
4. A system according to claim 3 comprising the steps of:-
 - a) extracting measles RNA and detecting the same using
20 NASBA,
 - b) enriching specific measles virus RNA/mRNA, and
 - c) repeating NASBA and/or RT-PCR.
5. A system according to any of claims 2 to 4 including a
25 buffered primer which primer comprises a 5' modified oligonucleotide sequence specific for the reverse transcribed DNA from an RNA measles virus.
6. A system according to any of claims 2 to 4 comprising a
30 RNA template specific for a RNA measles virus.
7. A system according to any of claims 2 to 5 wherein the primer is attached to a reporter molecule which enables post amplification detection of the DNA.
35
8. A system according to claim 7 wherein the reporter moiety is flurochromatic.

- 29 -

9. A system according to claim 5 comprising the following reagents:

- (1) M-MLV-reverse transcriptase;
- (2) random hexamers and/or oligo (dT)₁₂₋₁₈;
- 5 (3) a reaction buffer for (1) above;
- (4) a PCR reaction buffer; and
- (5) a 5' modified PCR primer with reporter molecules.

10. A system according to any preceding claim further
10 comprising a positive control comprising measles virus RNA in solution at a known concentration, and a negative control or means for providing the same.

11. A system according to any preceding claim wherein the
15 distinctive metabolic product including mRNA, protein or antigen is specifically distinctive of a wild type or an attenuated vaccine measles virus.

12. A system according to claim 11 wherein the distinctive
20 metabolic product is selected from a gene sequence or a metabolic product thereof to form a nucleocapsid protein, a phosphoprotein, a large protein, a matrix protein, a RNA polymerase complex, a haemagglutinin protein, a fusion protein and/or another characteristic product thereof detectable by
25 an immunological method.

13. A system according to claim 12 wherein the immunological
detection method is effected by immunohistochemistry or an ELISA assay, radioimmunoassay or enzyme linked gel assay.

30

14. A method for the analysis for Crohn's Disease and/or ulcerative colitis which method comprises the steps of:-

- (a) obtaining a tissue sample and securing the same in an enclosed reaction container;
- 35 (b) adding a reagent comprising a PCR buffer; MgCl₂, dNTP'S, random hexamers and diluting aqueously to a desired dilution factor;

- 30 -

(c) subsequently closing the enclosed reaction container, adding M-MLV-RT and subjecting to heat cycling for at least one cycle;

(d) treating with a washing buffer;

5 (e) adding a buffered Taq DNA polymerase, $MgCl_2$, DNTP and a primer at a predetermined dilution and then heat cycling for at least 25 cycles; and

(f) subsequently repeating step (d) and viewing for labelled product to indicate the presence of a measles virus RNA.

10

15. A method according to claim 14 wherein a Dig-11 or fluorescent labelled base primer is used.

16. A method for the in vitro analysis of a sample of Crohn's
15 Disease and/or ulcerative colitis which method comprises abstracting measles virus RNA from a suitable tissue and

(a) adding thereto a buffer further comprising DTT, DNTP's and RNase inhibitor and oligo (Dt) 12-18 and letting down the aqueous concentration in an enclosed reaction container;

20 (b) adding thereto M-MLV-RT and subjecting to incubation to provide a cDNA product;

(c) purifying the product and adding thereto a PCR buffer along with $MgCl_2$, dNTP's and an outer primer;

25 (d) subsequently adding Taq DNA polymerase and heating for at least 15 cycles;

(e) recovering an aliquot of a so-formed reaction production and adding buffered PCR with an inner primer and further Taq DNA polymers and recycling for at least 15 cycles; and

30 (f) removing the so-formed product and adding a loading dye and subjecting the dyed product to an electrophoresis to identify a selected product band.

(g) sequencing of the amplified products or hybridising the amplified products with a homologous or heterologous specific probe to distinguish vaccine strain measles virus from "wild"
35 type measles virus.

- 31 -

17. A method for the in vitro analysis of a sample of Crohn's Disease and/or ulcerative colitis which method comprises abstracting measles virus RNA from a suitable tissue and

(a) adding thereto a buffer further comprising UTP, and
5 primer mixes and warmed to room temperature and

(b) aliquoted into a reaction mixture comprising the template RNA, which mixture is heated for 5 min at 65°C, then at 41°C for 5 min. and

(c) subsequently adding the enzyme mix further comprising a
10 RNA polymerase and

(d) incubating at 41°C for 90 min in a water bath and

(e) detecting the reaction product by Northern blotting or an enzyme-linked gel assay.

(f) sequencing of the amplified products or hybridising the
15 amplified products with a homologous or heterologous specific probe to distinguish vaccine strain measles virus from "wild" type measles virus.

18. A system according to any preceding claim wherein the
20 annealing temperatures used during the amplification reaction are between 40°C and 65°C.

19. A system according to any preceding claim wherein the
method is capable of distinguishing between the attenuated
25 measles virus vaccine and the wild type virus.

20. A medicament for the treatment of Crohn's disease and/or ulcerative colitis which medicament comprises means for the prevention of expression, replication, transcription, RNA
30 processing and/or mRNA transport or any combination thereof of the measles virus *in vivo*.

21. A medicament as claimed in claim 19 comprising a measles virus antisense RNA for either genomic or antigenomic RNA and
35 a vector therefor.

- 32 -

22. A medicament as claimed in claim 19 wherein the medicament comprises a measles virus peptide or carbohydrate antigen or a monoclonal or polyclonal analogue thereof.

- 5 23. A measles vaccine comprising part or all of the measles virus genome encoded into an expression system adapted to induce a specific immune response to the measles virus and/or its component protein antigens.



Figure 1e



Figure 2c



Figure 1c



Figure 2b



Figure 1a




Figure 2a




Figure 4a



Figure 5b



Figure 3d



Figure 5a




Figure 3a




Figure 4b



Figure 1B







Figure 2D





Figure 5C and 5D (inset)



INTERNATIONAL SEARCH REPORT

Inter. Appl. No.
PCr/GB 96/00706

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12Q1/68 A61K39/165

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C12Q A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ANN MED, DEC 1993, 25 (6) P557-61, FINLAND, XP002009032 SMITH MS ET AL: "Viral association with Crohn's disease."	1,2
Y	see the whole document ---	3-19,21
X	J MED VIROL, APR 1993, 39 (4) P345-53, UNITED STATES, XP002009033 WAKEFIELD AJ ET AL: "Evidence of persistent measles virus infection in Crohn's disease."	1,2
Y	see the whole document ---	3-19,21
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

22 July 1996

Date of mailing of the international search report

07. 08. 96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Gurdjian, D

INTERNATIONAL SEARCH REPORT

International Application No

PCI/GB 96/00706

C(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	VIROLOGY, FEB 20 1995, 207 (1) P168-78, UNITED STATES, XP002009034 KOSCHEL K ET AL: "Measles virus antisense sequences specifically cure cells persistently infected with measles virus." see the whole document ---	21
X	J INFECT DIS, NOV 1994, 170 SUPPL 1 PS15-23, UNITED STATES, XP002009035 BELLINI WJ ET AL: "Virology of measles virus." see the whole document ---	23
Y	see the whole document ---	11-13,22
X	US,A,4 871 759 (MERCER JAMES B) 3 October 1989 see the whole document ---	20
Y	see the whole document ---	22
Y	LANCET, MAR 11 1995, 345 (8950) P660, ENGLAND, XP002009036 WAKEFIELD AJ ET AL: "Measles virus in Crohn's disease [letter;comment]" see the whole document -----	3-19

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 96/00706

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4871759	03-10-89	US-A- 3856966	24-12-74
		US-A- 4537775	27-08-85
		US-A- 3952103	20-04-76
		US-A- 4952594	28-08-90
		US-A- 5102902	07-04-92
		US-A- 4073928	14-02-78
		US-A- 4177281	04-12-79
